

Original Article

Evaluation of Root Dilaceration by Cone Beam Computed Tomography in Iranian South Subpopulation: Permanent Molars

Bahar Asheghi¹, DMD, MScD; Safoora Sahebi¹, DMD, MScD; Maryam Zangoeei Booshehri², DMD, MScD; Farnaz Sheybanifard³, DDS;

¹Dept. of Endodontics, School of Dentistry, Shiraz University of Medical Sciences, Shiraz, Iran.

²Dept. of Oral and Maxillofacial Radiology, School of Dentistry, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

³Student Research Committee, School of Dentistry, Shiraz University of Medical Sciences, Shiraz, Iran.

KEY WORDS

Cone-beam computed tomography (CBCT);
Dilacerations;
Molar teeth;

Received: 29 May 2021;
Revised: 22 August 2021;
Accepted: 12 October 2021;

Copyright

© Journal of Dentistry, This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International License, (<http://creativecommons.org/licenses/by/4.0/>) which permits reusers to copy and redistribute the material in any medium or format if the original work is properly cited, and attribution is given to the creator. The license also permits for commercial use.

ABSTRACT

Statement of the Problem: Root dilaceration is a developmental anomaly, which was first described in the 19th century as an abrupt change in the axial inclination between the crown and root of a tooth. Such an anomaly would potentially alter the dental therapeutic procedures and often demands special care. Hence, it is of great importance to diagnose dilaceration before starting any treatment procedure on the tooth.

Purpose: The aim of the current study was to evaluate the prevalence of dilaceration in the first and second maxillary and mandibular molar teeth in Shiraz, Iran, using cone-beam computed tomography (CBCT) radiography.

Materials and Method: In this retrospective study, 472 teeth from 141 CBCT images were studied. Each tooth was assessed by CBCT to diagnose dilaceration, its severity, direction, and anomaly location.

Results: In this study, among 472 studied teeth, 136 teeth (28.8%) were diagnosed with root dilaceration. The most frequently affected tooth was the maxillary second molar tooth (45%). The results showed a statistically significant relationship between the gender of the patients and the prevalence of dilaceration. Most dilacerations were observed in distal direction, with a mild intensity, and located in the apical third of the roots.

Conclusion: In this study, CBCT was presented as an advanced method for diagnosing dilaceration in maxillary second molar teeth and especially in the mesial root, which can be influenced by gender and it can play an important role in designing the treatment plans.

Corresponding Author: Sheybanifard F, School of Dentistry, Shiraz University of Medical Science, Shiraz, Iran. Tel: +98-7136263193-4 Email: sheybanifard.f@gmail.com

Cite this article as: Asheghi B, Sahebi S, Zangoeei Booshehri M, Sheybanifard F. Evaluation of Root Dilaceration by Cone Beam Computed Tomography in Iranian South Subpopulation: Permanent Molars. *J Dent Shiraz Univ Med Sci.*, 2022 September; 23(2 Suppl): 369-376.

Introduction

First introduced by Tomes [1] in 1848, dilaceration is an evolutionary disorder between the mineralized and non-mineralized parts of the structure of a developing tooth. This anomaly appears as a deviation of the axial inclination of the crown and roots of a tooth and can occur anywhere along the length of the tooth from the crown to the root apex [2-3].

Considering the inevitable presence of some degree of curvature in most tooth roots, some studies have defined dilaceration as a certain degree of curvature from

the normal dental longitudinal axis. For instance, Hamasha *et al.* [4] and Malcic *et al.* [5] considered dilaceration as the curvature of 90 degrees or more, whereas Chohayeb *et al.* [6] regarded the curvature of 20 degrees or more as dilaceration.

The development of dilaceration is associated with multiple parameters [7]. In this regard, it has been shown that the displacement of the root sheath of teeth following the gradient of bone remodeling within the alveolar bone can have a potential impact on the development of dilacerations [3]. Moreover, ethnicity and

gender are crucial parameters to consider. Therefore, the prevalence of dilaceration has been studied in multiple ethnic groups. It has had different frequencies ranging from 0.3% to 98% in the studied teeth [4-6, 8-9]. Another influential parameter in the development of dilaceration is trauma. It is worth noting that since dilaceration is very common in posterior teeth and trauma rarely affects this area, trauma was excluded from the list of influential parameters [4, 10]. Two other determinants of dilaceration in posterior teeth are the genetic alteration of the tooth germ position and the effect of delayed eruption of teeth [5, 9, 11-12]. Nevertheless, they do not seem to be the only reasons for dilaceration in these regions [3].

With respect to the location of dilaceration, some studies declare that dilaceration is more common in the posterior regions of the mandibular [4] and permanent teeth [13], while others state that the maxilla is more involved [5]. It has been shown that dilacerations in incisors, canines, and premolars is more common in the apical third of the roots. In molars, dilaceration usually occurs in the middle third of the roots; while in third molars, dilaceration occurs more frequently in the coronal third of the roots [5]. Nevertheless, there is no consensus among researchers on the teeth, which are the most, and the least affected with this anomaly [14].

Considering the above-mentioned parameters, it is of great importance to diagnose dilaceration at the roots. Radiography is the most efficient method for this purpose [4]. Most previous studies have used conventional radiographies such as periapical and panoramic to investigate the prevalence of dilacerations [15-18]. However, these methods can take the images of the teeth only in two dimensions.

An appropriate method to overcome this shortcoming is to use cone-beam computed tomography (CBCT) imaging. CBCT imaging is a technique that has been widely used in various studies of root and canal system morphology to show the third dimension of the teeth [19-20]. In this imaging technique, all the distortions and superimpositions are removed, making it more appealing in various teeth anomaly studies [21]. Therefore, using CBCT radiography leads to an accurate identification of the location, direction, and degree of dilacerations [21-22]. Nevertheless, only a few studies (case reports) have exclusively used CBCT imaging to

investigate the prevalence of dilacerations [23-24]. The aim of the current study was to use CBCT to investigate the prevalence of dilaceration in the teeth and roots of the first and second molars in the maxilla and mandible. Moreover, the severity, degree, and location of dilaceration in these teeth were studied. Furthermore, the relationship between the gender and the prevalence of dilaceration among patients referring to dental clinics in Shiraz, Iran, were investigated.

Materials and Method

Sample collection

In this cross-sectional and retrospective study, the prevalence of dilaceration was assessed in randomly selected high-quality CBCT radiographic images archived from 141 randomly selected patients (18 to 65 years old). These data were gathered from the archives of four oral and maxillofacial radiology clinics in Shiraz from July 2017 to August 2020. The CBCT images were taken from patients with different conditions such as dental implantation, maxillary and mandibular bone fracture, or tumor. Therefore, additional radiation was avoided.

The current study was approved by the Ethics Committee of Shiraz University of Medical Sciences (IR.SUMS.DENTAL.REC.1398.33).

In this study, the presence of dilaceration in 141 CBCT images, 472 maxillary and mandibular molars, and 1204 molar roots was evaluated. Of all molars, 119 were maxillary first molars, 140 were maxillary second molars, 95 were mandibular first molars, and 118 were mandibular second molars.

In the present study, high-resolution CBCT images of the first and second mandibular and maxillary molars were selected. All images were taken from completely erupted teeth with closed and fully developed apices.

Furthermore, the supernumerary teeth, teeth with internal and external resorption, and deciduous or permanent teeth with open apex roots were excluded from the study.

The data about the number of tooth with dilaceration, the patients' gender, the position of the teeth with dilaceration in the jaw (maxilla or mandible), the presence or absence of dilaceration in each root (categorized based on the type of root), the severity of dilaceration based on Santana's classification [25], the direction of dilaceration (mesial, distal, buccal, palatal/lingual), the

location of dilaceration (apical, middle, coronal), the teeth with S-shaped roots, and the presence of supernumerary roots in the molar teeth and dilaceration in them were collected for statistical analysis.

In the present study, the definition of dilaceration presented by Chohayeb *et al.* [6] was used. They consider the deviations of more than 20 degrees from the longitudinal axis of the tooth as dilaceration. Moreover, according to the study of Santana [25], the intensity of the curvature was categorized as mild (with 20-40 degrees of deviation), moderate (with 41-60 degrees of deviation), and severe (with more than 60 degrees of deviation).

The bending angle of dilaceration was investigated using the method presented by Schneider [26]. In this approach, the angle between two hypothetical lines connecting the apex and the orifice to the beginning of the curvature is measured.

In the current study, the presence of at least one root with dilaceration in the multi-rooted teeth was considered enough to mark that tooth as dilacerated. Furthermore, in teeth with multiple dilacerations in different planes, the largest angle was considered as dilaceration.

All CBCT images were prepared using a Planmeca Promax 3D Mid device (Helsinki, Finland) at 90 kVp and 14 Ma, with an exposure time of 15 s, and automatically adjusted based on the patients' body weight and size. A maximum field of view of 10*10 cm, a voxel size of 150 μ , and a high definition mode were used.

Root dilaceration was evaluated using a magnification tool in the Romexis software. The CBCT images in the axial, coronal, and sagittal sections were analyzed using the Romexis imaging software (version: 3.8.2) on a 32-inch monitor in dim light (Figure 1).

All CBCT images were inspected retrospectively by a calibrated endodontist and a senior dental student who were trained to interpret the images independently. There was a two-week interval between the assessments. Prior to the experiments, the investigators evaluated 60 other CBCT images. If any disparity was seen between the opinions, a radiologist evaluated the images in order to reach a consensus. To assess the intra-examiner reliability, a re-assessment was performed one month after the first session.

Data analysis

The data in the current study were analyzed by the SPSS software (version: 18.0, SPSS Inc., Chicago IL, USA) and the figures were generated using the Graph-Pad Prism software (version: 8.0). The Chi-square statistical test and Fisher's exact test were used to evaluate the qualitative data and the significance level was set at $p < 0.05$. In this study, the p value was used to investigate whether root dilaceration was statistically dependent on gender, jaw type, tooth type, and root type. The prevalence of dilaceration, S-shaped roots, and extra roots in the studied teeth was presented as percentages. The agreements between the inter- and intra-examiners were calculated using Cohen's kappa coefficient.

Results

It was observed that the overall prevalence of dilaceration in at least one root in the studied population was 51.7% ($n=73$). The prevalence of dilaceration in the first and second molars was 28.8% ($n=136$) (Figure 2). The assessments of the relationship between the prevalence of dilaceration and the patients' gender revealed that dilaceration was statistically dependent on gender and occurred more often in females ($p=0.044$) (Table 1).




Figure 1: Moderate dilaceration in the mesial root of maxillary second molar in the sagittal (left) and 3D (right) planes




Figure 2: The prevalence of dilaceration in the studied teeth

Table 1: The distribution of dilaceration between the two genders

	Female	Male	Total	<i>p</i> Value
Teeth without dilaceration	186 (67.6%)	150 (76.1%)	336 (71.2%)	<i>p</i> = 0.044
Teeth with dilaceration	89 (32.4%)	47 (23.9%)	136 (28.8%)	
Total	275 (100%)	197 (100%)	472 (100%)	

Table 2: The prevalence and distribution of the dilacerated teeth based on the jaws

	Maxilla	Mandible	Total	<i>p</i> Value
Teeth without dilaceration	160 (61.8%)	176 (82.6%)	336 (71.2%)	<i>p</i> < 0.001
Teeth with dilaceration	99 (38.2%)	37 (17.4%)	136 (28.8%)	
Total	275 (100%)	213 (100%)	472 (100%)	

The results obtained in this study demonstrated that the prevalence of dilaceration was more in the upper jaw than in the lower jaw ($p < 0.001$) (Table 2). The teeth most affected with this anomaly were the maxillary second molars ($p < 0.001$).

The maxillary first molars, the mandibular first mol-

lars, and the mandibular second molars respectively occupied the next ranks (Figure 3a).

Investigating the roots of the above-mentioned teeth showed that dilaceration was more common in the mesial roots of maxillary second molars ($p < 0.001$) (Figure 3b). In general, it was shown that curvature and root dilaceration were more common in the distal direction mostly in the third apical with a mild intensity ($p < 0.001$) (Figure 4).

Out of 472 teeth, 2.1% had at most two extra roots in which the prevalence of dilaceration was 12.5% ($n=1$). Moreover, it was observed that 5.72% of all the teeth had S-shaped roots.

Regarding the inter-examiner's agreement (after the training session), Cohen's kappa coefficients for the first and second assessments were 0.991 and 0.993, respectively. The overall Cohen's kappa coefficient for the intra-examiner's agreement was 0.996. Taken together, this result suggests a very good agreement between the intra- and inter-examiners.

Discussion

Dilaceration is recognized as one of the most important anomalies that have the potential to affect the success rate of endodontic treatments [4, 6, 27]. Frequent errors such as ledge, transport, zipping, and broken file are the results of abnormal root canal curvature, which in turn can affect the outcome of endodontic treatment procedures especially in molar teeth [28]. Hence, it is of great importance to diagnose dilaceration before starting any treatment procedure on the tooth. In severe cases of dilaceration, alternative endodontic treatment methods




Figure 3: a) The prevalence of dilaceration in each studied tooth, b) based on the root positions




Figure 4: a: The direction, b: Everity, c: Prevalence of dilaceration in the root length

such as vital pulp therapy should be considered [4, 6].

CBCT has been introduced in the current study as a novel, non-invasive, and accurate radiographic technique for the diagnosis of dilaceration. It enables dentists to examine the tooth in three dimensions. This method has some advantages over invasive techniques such as tooth extraction by providing the exact angle of dilaceration at the root. Moreover, CBCT imaging provides some information about the patient's age, gender, and teeth symmetry [29-31].

In the current study, the different planes of the roots of the first and second molar teeth were investigated using CBCT images. Such planes are not visible in conventional radiographs. Here, the criteria presented by Chohayeb [6] and the classification of Santana *et al.* [25] were used to define dilaceration based on root curvature. These criteria (deviation of >20) were also used in different thirds of the roots.

The literature review shows that there is a large discrepancy in the prevalence of dilaceration in different populations. It has previously been shown that dilaceration is the most common dental anomaly in the population of Iran, comprising 15% of all types of dental anomalies [32]. Due to the vastness of the country, few studies have investigated the prevalence of dilaceration in Iran and most of them have been conducted without CBCT radiography [17-18, 32].

In the current study, the prevalence of dilaceration was reported among patients diagnosed with at least one case of this anomaly (51.7%). The teeth that had at least one root with dilaceration (28.8%) were also investigated.

Multiple studies assessed the prevalence of dilaceration in molar teeth using periapical or panoramic radiographies. They showed that the prevalence of dilaceration was below 9% [5, 9, 14-15, 18]. A higher prevalence of dilaceration in molar teeth was observed in the present study (28.8%) than in previous studies. This may be due

to the better diagnosis of buccal and/or lingual dimensions via CBCT imaging. Moreover, different sample sizes, different races, and different definitions of dilaceration were used in the current study, potentially affecting the final results.

In line with some previous studies [9, 15, 33], the current study showed that the presence of dilaceration was significantly dependent on the patients' gender and was more common in females. However, some other papers showed that dilaceration was more common in males [32, 34] or was even gender-independent [14, 16, 35]. It was observed that most teeth affected by dilaceration were the maxillary second molars. Irrespective of the third molars, a similar observation has been reported in two previous studies [16, 36]. Meanwhile, excluding the third molars, some studies have shown a different result and have introduced other teeth such as maxillary first molars [5,14] or mandibular second molars [9, 15, 18] as the candidates with a higher prevalence of dilaceration. There are controversies among different studies regarding the location of dilaceration; however, two studies, which considered all teeth in their investigations, have mentioned the posterior region as the most common location of dilacerations [9, 16].

In the current study, it was also observed that the prevalence of dilaceration was significantly higher in the maxilla than in the mandible as has been previously reported [5,9,36]. Although some studies have indicated that dilaceration mostly occurs in the mandible [4, 15], previous investigations have not reported a difference between the two jaws [14, 16, 18, 33].

Inspecting the roots separately showed that the highest prevalence of dilaceration was in the mesial root of the maxillary second molar. Nevertheless, the distal root of the mandibular second molar has also been introduced as a candidate with the greatest prevalence of dilacerations [18]. In the present study, the extra roots of

mandibular molars as well as the buccal roots of maxillary second molars with two roots were also reported. Mandibular molar roots are referred to as entomolaris and paramolaris corresponding to teeth numbers 6 and 7, respectively [1,18]. The prevalence of dilaceration in the aforementioned roots is shown in Figure 3B. To the best of our knowledge, this was the first study to evaluate the prevalence of dilaceration in these teeth using CBCT radiography.

A dilaceration with a mild intensity was the most common dilaceration in the current study. This was consistent with the studies of Santana *et al.* [25] and Erlich and Panella [37]. This was followed by moderate and severe intensities, respectively. Furthermore, examining the location of dilaceration on the roots of the teeth revealed that the highest prevalence was related to the third apical of the roots. This is in line with some previous studies [5, 25, 36, 37-38] and in contrast with another research, which showed that dilaceration was more common in molars in the middle third [5].

The results of the present study confirmed those of previous studies and showed that dilaceration was mainly in the distal direction [25, 34, 39].

Due to the relatively high prevalence of dilaceration in this study and the vastness of Iran, these epidemiological data cannot be generalized to the whole country. Therefore, it is recommended that similar studies with larger sample sizes and in different parts of the country be conducted using CBCT radiography.

Conclusion

Using an advanced imaging modality, the results showed that dilaceration was relatively common in the molar teeth and especially in the mesial root, which can be influenced by gender and the majority of participants suffered from some degree of curvature. Thus, it is highly recommended that dental practitioners evaluate the presence of dilaceration with the help of diagnostic imaging before starting any treatment on molar teeth. This is even more crucial when the patient is female or the teeth are maxillary second molars.

Acknowledgments

The Ethics Committee of Shiraz University of Medical Sciences approved the current study (IR.SUMS. DENTAL.REC.1398.33). All methods were carried out in

accordance with the institutional guidelines and regulations. The signed informed consents were obtained from the patients so that their radiographic images could be used in the current study.

Conflict of Interests

The authors declare that they have no conflicts of interests.

References

- [1] Tomes J. A Course of Lectures on Dental Physiology and Surgery, delivered at the Middlesex Hospital School. *Am J Dent Sci.* 1848; 8: 209.
- [2] Institution BS. British standard glossary of dental terms. Available at: <https://iucatl.iu.edu/iupui/2203278>
- [3] Standerwick R. A possible etiology for the dilaceration and flexion of permanent tooth roots relative to bone remodeling gradients in alveolar bone. *Dent Hypotheses.* 2014; 5: 7-10.
- [4] Hamasha AA, Al-Khateeb T, Darwazeh A. Prevalence of dilaceration in Jordanian adults. *Int Endod J.* 2002; 35: 910-912.
- [5] Malčić A, Jukić S, Brzović V, Miletić I, Pelivan I, Anić I. Prevalence of root dilaceration in adult dental patients in Croatia. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2006; 102: 104-109.
- [6] Chohayeb AA. Dilaceration of permanent upper lateral incisors: frequency, direction, and endodontic treatment implications. *Oral Surg Oral Med Oral Pathol.* 1983; 55: 519-520.
- [7] Jafarzadeh H, Abbott PV. Dilaceration: review of an endodontic challenge. *J Endod.* 2007; 33: 1025-1030.
- [8] Miloglu O, Cakici F, Caglayan F, Yilmaz AB, Demirkaya F. The prevalence of root dilacerations in a Turkish population. *Med Oral Patol Oral Cir Bucal.* 2010; 15: e441-e444.
- [9] Udoye CI, Jafarzadeh H. Dilaceration among Nigerians: prevalence, distribution, and its relationship with trauma. *Endod Dent Traumatol.* 2009; 25: 439-441.
- [10] White SC, Paroah MJ. White's and Pharoah's Oral Radiology: Principles and Interpretation. 8th ed. CV Mosby Co: St Louis; 2019. p.817.
- [11] Proffit WR, Fields HW, Larson B, Sarver DM. Contemporary orthodontics-e-book. Elsevier Health Sciences: St Louis; 2018. p.130.
- [12] Regezi JA, Sciubba J, Jordan RC. Oral pathology: clinic-

- al pathologic correlations. 7th ed. Elsevier Health Sciences: St Louis; 2016. p. 375.
- [13] Tiecke RW, Stuteville OH, Calandra JC. Pathologic physiology of oral disease. CV Mosby Co: St Louis; 1959. p. 451.
- [14] Miloglu O, Cakici F, Caglayan F, Yilmaz AB, Demirkaya F. The prevalence of root dilacerations in a Turkish population. *Med Oral Patol Oral Cir Bucal*. 2010; 15: e441-e444.
- [15] Colak H, Bayraktar Y, Hamidi MM, Tan E, Colak T. Prevalence of root dilacerations in Central Anatolian Turkish dental patients. *West Indian Med J*. 2012; 61: 635-639.
- [16] Karadaş M, Tahan E. Prevalence of Root Dilaceration in a Subpopulation of Northeast Turkey. *Türkiye Klinikleri Dishekimligi Bilimleri Dergisi*. 2015; 21: 71-76.
- [17] Kuzekanani M, Sadeghi MT. Prevalence and distribution of dilacerations in the permanent dentition of an Iranian population. *Eur J Anat*. 2019; 23: 273-277.
- [18] Nabavizadeh M, Shamsi MS, Moazami F, Abbaszadegan A. Prevalence of root dilaceration in adult patients referred to Shiraz dental school (2005-2010). *J Dent (Shiraz)*. 2013; 14: 160.
- [19] Asheghi B, Sahebi S. Morphological evaluation of maxillary premolar canals in Iranian population: a cone-beam computed tomography study. *J Dent (Shiraz)*. 2020; 21: 215.
- [20] Herrero-Hernández S, López-Valverde N, Bravo M, Valencia de Pablo Ó, Peix-Sánchez M, Flores-Fraile J, et al. Root canal morphology of the permanent mandibular incisors by cone beam computed tomography: a systematic review. *Appl Sci (Basel)*. 2020; 10: 4914.
- [21] Agnihotri A, Marwah N, Dutta S. Dilacerated unerupted central incisor: A case report. *J Indian Soc Pedod Prev Dent*. 2006; 24: 152.
- [22] Crescini A, Doldo T. Dilaceration and angulation in upper incisors consequent to dental injuries in the primary dentition: orthodontic management. *Prog Orthod*. 2002; 3: 29-41.
- [23] Bletska A, Bessonova MV, Iden O. Using cone-beam CT in diagnosis and management of severe dilaceration following trauma in primary teeth: a case report. *Int J Clin Pediatr Dent*. 2020; 13(Suppl 1): S110.
- [24] Mahesh R, Kanimozhi I, Sivakumar M. Dilaceration and eruption disturbances in permanent teeth: a sequelae of trauma to their predecessors-diagnosis and treatment using Cone Beam CT. *J Clin Diagn Res*. 2014; 8: ZD10.
- [25] Santana EJBd CA, Tavano O. Determinação da prevalência e estudo morfológico da dilaceração radicular. *Rev Fac Odontol Univ Fed Bahia*. 1993; 12: 40-52.
- [26] Schneider SW. A comparison of canal preparations in straight and curved root canals. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 1971; 32: 271-275.
- [27] Neville BW, Damm DD, Allen C, Chi AC. Oral and maxillofacial pathology. 4th ed. Elsevier Health Sciences: WB Saunders; 2015. p. 604-605.
- [28] Neelakantan P, Subbarao C, Ahuja R, Subbarao CV. Root and canal morphology of Indian maxillary premolars by a modified root canal staining technique. *Odontology*. 2011; 99: 18-21.
- [29] Matherne RP, Angelopoulos C, Kulild JC, Tira D. Use of cone-beam computed tomography to identify root canal systems *in vitro*. *J Endod*. 2008; 34: 87-89.
- [30] Nair MK, Nair UP. Digital and advanced imaging in endodontics: a review. *J Endod*. 2007; 33: 1-6.
- [31] Neelakantan P, Subbarao C, Subbarao CV. Comparative evaluation of modified canal staining and clearing technique, cone-beam computed tomography, peripheral quantitative computed tomography, spiral computed tomography, and plain and contrast medium-enhanced digital radiography in studying root canal morphology. *J Endod*. 2010; 36: 1547-1551.
- [32] Ardakani FE, Sheikha M, Ahmadi H. Prevalence of dental developmental anomalies: a radiographic study. *Community Dent Health*. 2007; 24: 140.
- [33] Ledesma-Montes C, Hernández-Guerrero JC, Jiménez-Farfán MD. Frequency of dilaceration in a Mexican school-based population. *J Clin Exper Dent*. 2018; 10: e665.
- [34] Silva BFd, Costa LED, Beltrão RV, Rodrigues TL, Farias RLd, Beltrão RTS. Prevalence assessment of root dilaceration in permanent incisors. *Dental Press J Orthod*. 2012; 17: 97-102.
- [35] Fuentes R, Farfán C, Astete N, Navarro P, Arias A. Distal root curvatures in mandibular molars: analysis using digital panoramic X-rays. *Folia Morphologica*. 2018; 77: 131-137.
- [36] Karataş E, Topçuoğlu HS, Arslan H, Erdoğan Ş, Ezmeci EB. Prevalence of root dilacerations in an east anatolian endodontic patient population. *Türkiye Klinikleri Diş Hekimliği Bilimleri Dergisi*. 2012; 18: 28-32.
- [37] Erlich T FPM, Panella J. Estudo da prevalência da dilace-

ração radicular, por meio de exame radiográfico periapical, numa amostra populacional da Grande São Paulo. *RPG Rev Pos-Grad.* 2001; 8(abr./ju 2001): 129-137.

[38] Estrela C, Bueno MR, Barletta FB, Guedes OA, Porto OC, Estrela CR, et al. Identification of apical and cervical curvature radius of human molars. *Braz Dent J.* 2015; 26:

351-356.

[39] Aminsobhani M, Sadegh M, Meraji N, Razmi H, Kharazifard MJ. Evaluation of the root and canal morphology of mandibular permanent anterior teeth in an Iranian population by cone-beam computed tomography. *J Dent (Tehran).* 2013; 10: 358.