

Original Article

The Knowledge and Attitude of Diabetic Patients Regarding Oral and Dental Disorders in Kerman Diabetes Clinics

Shahla Kakoei, DDS, MSc¹; Salimeh Afzali, DDS²; Masoud Parirokh, DDS, MSc³; Sina Kakoei, DVM, PhD⁴, Mahshid Mostafavi, DVM, PhD⁵, Amir Nekouei, MSc³;

¹ Neuroscience Research Center, Kerman University of Medical Sciences, Oral Medicine Dep. Kerman, Iran.

² Oral and Dental Diseases Research center, Kerman University of Medical Sciences, Kerman, Iran.

³ Endodontology Research Center, Kerman University of Medical Sciences, Kerman, Iran.

⁴ Physiology Research Center, Kerman University of Medical Sciences, Ph.D. candidate of Endodontology, Kerman, Iran

⁵ Leishmaniasis Research Center, Kerman University of Medical Sciences, Kerman, Iran.

KEY WORDS

Oral disease;
Diabetes;
Knowledge;
Attitude;

ABSTRACT

Statement of the Problem: Diabetes mellitus (DM) is one of the most common endocrine disorder. This disease has devastating effects on many organs and tissues of the body including heart, kidney, mouth, teeth and etc.

Purpose: The aim of this study was to evaluate the knowledge and attitude of diabetic patients about dental and oral diseases.

Materials and Method: In this cross-sectional study, 433 diabetic patients who referred to Kerman diabetes clinics, were included. Data were collected using a questionnaire consisted of three parts of demographic characteristics, knowledge of oral and systemic complications of DM, and patients' attitude regarding their oral health. Data were analyzed using t-test and multiple linear regression test by SPSS version 21. Statistically significant values were considered at $p \leq 0.05$.

Results: The mean scores for the knowledge of systemic and oral complications were 0.80 ± 0.21 and 0.39 ± 0.23 , respectively. The mean total knowledge of diabetic patients was 0.53 ± 0.18 , and the mean scores for the patients' attitude was 0.63 ± 0.11 . It was revealed that people with a family history of diabetes did not have significantly greater overall knowledge ($p = 0.082$). Also, people with longer disease duration ($p = 0.004$) and female patients ($p = 0.05$) had significantly a better knowledge and attitude in terms of oral health.

Conclusion: The knowledge and attitude of patients regarding their oral and dental health and diseases were at moderate level, which should be promoted by constant planning and education according to the current needs of society.

Received:
Revised:
Accepted:

Corresponding Author: Kakoei S, Physiology Research Center, Kerman University of Medical Sciences, Kerman, Iran. Area code 76169-14115 Email: s.kakoei@yahoo.com Tel: +98-34-32264071 Fax: +98-34-32264097

Cite this article as:

Introduction

With increasing age of patients referring to dental clinics, dentists may face more with patients who suffer from some diseases. In a study conducted in three cities of Iran, about 50% of the patients had at least one systemic disease [1]. Diabetes mellitus (DM), which is sometimes known as silent epidemic, is one of the important chronic metabolic diseases and a major health problem worldwide. The prevalence of this disease is increasing especially in developing countries [2]. Cur-

rently, diabetes is a major health problem in all developed and developing countries, and its prevalence is increasing day by day in various countries, including Iran. According to the World Health Organization (WHO) report, the prevalence of type 2 diabetes in Iran was 5.7% in 2000 and it will be 6.8% in 2025. Accordingly, the diabetic population in these years will be 1,977,000 and 5,125,000, respectively [3]. Type 2 diabetes imposes a heavy and increasing burden on medical care all across the world [3]. The prevalence of dia-

betes increased from 30 million in 1985 to 135 million in 1995, and it is estimated to be 366 million cases in 2030 [4]. Also, many of the new cases of diabetes are observed in developing countries, and it seems that the Middle East will have the highest increase in the prevalence of diabetes in 2030. Furthermore, it is estimated that in 2025, more than 75% of all diabetic patients will be in developing countries. In Iran, more than 3,000,000 people have diabetes, and according to WHO estimation, it will be nearly 7,000,000 cases in 2030. With a prevalence of more than 7%, Iran is amongst the areas with the highest prevalence of diabetes in the world. Annually, an average number of 500 thousand individuals will be added to diabetic population of Iran [5-6].

DM is considered as a metabolic disease with many complications such as cardiovascular, neurological, kidney, eye, nerve and mouth complications, and one of the major global health problems [7]. In addition to its effects on other organs, it can also affect oral health [8]. Oral symptoms of DM include dry mouth, burning mouth, taste disorders, large thyroid gland, periodontal disease, and bacterial and fungal infections [9-10]. Studies have shown that irreversible complications of diabetes are caused by glycation end products that make changes in cholesterol, albumin, collagen, and hemoglobin, providing the grounds for complications in diabetic patients [11]. Frequent urination with neurological and pathological changes in salivary glands reduces the amount of saliva in these patients. The oral symptoms of diabetes include decreasing saliva PH, dental caries, gingivitis, periodontitis, mouth irritation, dry mouth, changes in the chemical composition of saliva, oral yeast infections, median rhomboid glossitis (MRG), and oral lichen planus (OLP) [12].

Increasing information about the potential of periodontal diseases and xerostomia can be important for the prevention of oral problems in diabetic patients. Studies have also shown that increased information about proper oral health behaviors is necessary to take care of one's mouth [13]. Considering the prevalence of diabetes (7%) in Iran and its importance, as well as the existing oral diseases, in this study, the knowledge and attitude of dental and oral diseases in diabetic patients in Kerman, Iran.

Materials and Method

In this cross-sectional study, diabetic patients who ref-

erred to diabetes clinics in two Hospitals of Kerman (Shahid Bahonar and Afzalipour) were included. The participants were selected using simple random sampling method. The sample size was calculated using G*Power software version 3.0.1. Based on $\alpha = 0.05$, effect size = 0.50, and power = 80%, 441 participants were found to be sufficient.

The inclusion criteria were considered as: age between 17 and 75 years, HbA1C $\geq 6.5\%$ or fasting plasma glucose (FPG) ≥ 126 mg/dl or 2-hour plasma glucose ≥ 200 mg/dl, symptoms of hyperglycemia and hyperglycemia with a random plasma glucose ≥ 200 mg/dl [14], without mental problems.

The exclusion criteria were including of: age below 15 and over 75 years. HbA1C < 6.5 or FPG < 126 or 2-hour plasma glucose or random plasma glucose < 200 mg/dl or having mental problem.

This study was designed to measure the knowledge and attitude of oral diseases in diabetic patients. Data were collected using a three-part questionnaire: The first part consisted the demographic and individual characteristics such as age, sex, duration of diabetes, type of treatment, the condition of diabetes, family history of the disease, type of diabetes, and the most recent amount of HbA1c and FBS. The second part consisted questions related to knowledge in two areas: Knowledge about systemic diseases and oral complications related to DM. The second part included questions related to the patients' attitude regarding the prevention of oral and dental diseases such as brushing, flossing, and visiting dentists. The responses were based on three options: Yes, No, and I do not know.

The questionnaire content validity was approved by 10 experts of Kerman Dental School. The questions' intelligibility was discussed as well. Based on the experts' opinion, 19 questions were considered appropriate and very appropriate. Validity and reliability of the questionnaire were measured by content validity index (0.78), and Cronbach's alpha (0.80) was at acceptable level. The answers of questions were scored from 1 to 0 (true answers: 1, false answers: -1, and I don't know: 0). At the end, the knowledge of participants was scored as: good (0.67, 1), moderate (-0.67, 0.67), and poor (-1, -0.67).

An informed consent was obtained from all participants. The participants were also assured that their in-

formation would remain confidential and would be investigated for purely statistical reasons. In addition, for semi-literate and illiterate patients, the questions were read to them. The study was approved by the Ethics Committee of Kerman University of Medical Sciences (Ethical code: EC/KMRC/92-52).

The results were expressed as percentages and 95% confidence intervals. The data were analyzed using t-test and multiple linear regressions with significance level of 0.05 by SPSS version 21.

Results

In this study, 433 questionnaires were randomly distributed among selected diabetic patients, of which 417 questionnaires were returned. The mean age of patients was 52.1±12.1 years, while 66.5% of them were female and 53.8% had uncontrolled diabetes. At least, 85% of the participants have knowledge about the kidney and eye diseases, paresthesia, and delayed wound healing (Table 1). The participants’ knowledge about the oral complications is presented in Table 2. More than 90% of the patients mentioned xerostomia as the most frequent complication (Table 2).

Table 1: Knowledge questions about systemic complications of DM

Type of disease	Yes No(%)	No No(%)	I don't know No (%)
Kidney disease	386(92.5)	25(6)	6(1.4)
Eye disease	397(95.2)	15(3.6)	5(1.2)
Neurological disease	216(51.7)	121(29.1)	80(19.2)
Heart disease	326(78.3)	63(15)	28(6.8)
Mental disease	286(68.8)	77(18.4)	54(13)
Delayed wound healing	356(85.3)	47(11.8)	12(2.9)
Paresthesia	376(91.1)	26(6.3)	15(3.6)

Table 2: Knowledge questions about oral complications of DM

Oral Complications	Yes No(%)	No No(%)	I don't know No(%)
Halitosis	172(41.3)	160(38.4)	85(20.3)
Tooth discoloration	114(27.3)	184(44.1)	119(28.6)
Tooth sensitivity	135(32.4)	170(40.8)	112(26.8)
Tooth decay	219(52.5)	119(28.5)	79(19)
Tooth mobility	166(39.7)	162(38.9)	89(21.4)
Periodontitis	134(34.4)	190(54.4)	84(20.2)
Burning mouth sensation	140(3.7)	216(51.7)	61(14.6)
Aphthous ulcer	107(25.5)	251(60.2)	59(14.3)
Xerostomia	376(90.1)	34(8.2)	7(1.7)
Taste change	228(54.8)	137(32.8)	52(12.4)
Abscess	109(26.2)	240(57.5)	68(16.3)
Candidiasis	110(26.4)	246(59.1)	61(14.5)

Also, 64.1% of the participants (n=267) believed that blood sugar regulation can prevent oral diseases. In addition, 75% (n=313) of the participants believed that regular dental checkups can prevent the progression or incidence of oral diseases. Also, 46.1% (n=192) stated that they used a toothbrush or dental floss regularly, which can improve their oral health (Table 3).

The results of statistical analyses for questions related to dental and oral health attitude are presented in Table 4. About 60.5% of the participants brushed their teeth twice a day. Surprisingly, 66.8% of the participants did not use dental floss. In addition, more than 50% of them did not check up their teeth annually (Table 4).

The mean score for knowledge about systemic complications was 0.80±0.21, and 4.9% of the patients had poor knowledge about systemic complications, while 15.1% had moderate and 82% had good knowledge. The mean score of knowledge of oral complications was 0.39±0.23, and 50% of the patients had poor knowledge about oral complications, while 35.1% had moderate and 14.9% had good knowledge. The mean score of attitude was 0.63±0.11.

Table 3: Knowledge questions about oral disease prevention

Oral disease prevention	Yes No (%)	No No (%)	I don't know No (%)
Can blood sugar regulation prevent oral diseases?	267(64.1)	25(6)	125(29.9)
Can regular dental checkups prevent the progression or occurrence of oral diseases?	313(75)	18(4.4)	85(20.6)
Does using a toothbrush or dental floss on a regular basis improved oral diseases?	192(46.1)	74(17.7)	151(36.2)

Table 4: Attitude questions about dental and oral health

Practice questions	Answers	No.(%)
How many times do you brush daily?	Once	36(8.8)
	Twice	255(60.5)
	Three times	94(22.9)
Do you use Dental floss?	No	279(66.8)
	Some times	70(16.8)
	Yes	68(16.3)
If yes how many times?	Daily	69(16.5)
	Weekly	8(19)
	Monthly	40(9.6)
Do you visit dentist annually?	Yes	162(39)
	No	255(61)
If yes how many times?	One month	1(0.2)
	Three month	3(0.7)
	Six month	21(12.2)
	Annual	105(25.2)

The mean score was higher in female patients, and there were statistically significant differences in the knowledge of systemic complications ($p= 0.003$) and oral complications ($p= 0.037$) between the two genders. There was no significant difference between total score of knowledge and attitude in both genders. The age variable analysis showed that older people had significantly more knowledge about systemic complications. However, there was no significant differences between all of the dependent variables and familial history. But there was a significant relationship between disease duration and knowledge about systemic complications and total score of knowledge and attitude (Table 5).

Discussion

DM, as the most common metabolic disease, has wide-spread effects on body including mouth and teeth. The high incidence of candidiasis, periodontal disease, and dental decay in people with diabetes is also one of these effects. This is mainly due to the effects of disorder on gingival crevicular fluid (GCF) especially saliva [2,10]. The recent studies have shown that there is a relationship between increasing salivary immunoglobulin A (IgA) in diabetic patients and denture stomatitis and xerostomia [15-16]. It is reported that in 2000, 2.8% of people suffered from diabetes, and estimations show that this will amount to 4.4% in 2030 [17]. According to the latest statistics, there were nearly 4 million diabetic patients in Iran. According to the international statistics, this number will become three times more every 15 years [18]. Poor control of blood glucose due to diabetes leads to nephropathy, retinopathy, stroke, and coronary artery diseases. In addition, studies have shown that diabetic patients are 2 to 3 times more prone to periodontal diseases compared to healthy people. Several studies have proven a reciprocal relationship between periodontal disease and DM. Uncontrolled DM can cause periodontitis but surprisingly, treatment of periodontitis can improve blood sugar control [16].

In this study, the mean age of the patients was 52 ± 12 years, which was close to that reported by Yuen *et al.* [19] (57.9 ± 12.8 years). In this study, 66.5% of the patients were women that is similar to the gender distribution in the study of Delvarianzadeh *et al.* [20]. The mean duration of disease since diagnosis in patients was 8.8 ± 6.4 years. As age increases, the incidence of complications of the disease also increases [21]. Another study has reported that the mean duration of diabetes complications is 10.5 years [22]. Consistent with the results reported by Eldarrate *et al.* [23] and Allen *et al.* [24], the results of this study showed that diabetic patients' knowledge of other systemic diseases associated with diabetes is more than their knowledge of oral disease. The mean score for knowledge of systemic and oral complications was $80.4\pm 21.4\%$ and $39\pm 23.3\%$, respectively. The percentage of participants who were aware of increased risk of eye, kidney, and heart diseases was much more compared to the knowledge of gum disease, tooth decay, and fungal infections.

According to the results, the diabetic patients' knowledge of the possibility of developing oral diseases such as gum disease, tooth decay, and fungal infection caused by dry mouth was very inadequate. The results of this study showed that 50% of the participants had a low knowledge of the oral complications of diabetes. Several studies have reported that diabetic patients' knowledge of oral health and oral complications of diabetes is inadequate [25-27]. In this study, 90.1% of the participants were aware of the relationship between diabetes and dry mouth. Kakoei *et al.* [28] showed that xerostomia had most important factor on oral health impact profile (OHIP) in diabetic patients. In a study by Eldarrate, it was shown that more than 70% of the participants suffered from dry mouth while they were not aware of the harmful effects of dry mouth on health [23]. Yuen *et al.* [19] showed that only 30% of diabetic patients were aware of the effects of dry mouth on health.

Table 5: Relationship between knowledge and attitude with demographic variables

Dependent Variable independent Variable	Knowledge about systematic complications		Knowledge about oral complications		Total score of knowledge		Total score of attitude		Total score of Knowledge & attitude	
	p Value	β	p Value	β	p Value	β	p Value	β	p Value	β
Age	0.015	0.256	0.419	0.097	0.694	-0.108	0.082	-0.07	0.921	0.014
Sex	0.003	7.158	0.037	5.662	0.487	4.737	0.284	21.549	0.052	6.4
Familial history	0.534	-1.570	0.103	4720	0.082	12.05	0.254	-36.579	0.148	4.678
Duration of disease	0.048	0.388	0.355	0.210	0.928	0.049	0.699	0.362	0.004	0.930

Only 64% of the participants in this study believed that the blood sugar regulation can prevent oral diseases. Insufficient knowledge of the relationship between glycemic control in patients with diabetes and periodontal disease was remarkable in this study. It is clear that studies on the knowledge of the relationship between diabetes and periodontal disease in diabetic patients are not sufficient. It seems that the knowledge of such relationships should be strengthened.

In this study, only 46% of the patients believed that brushing or flossing can improve the disease treatment process. Also, it was revealed that follow-up treatments of periodontal diseases significantly improved blood sugar control in patients with type 2 diabetes [29].

In the present study, 52.9 % of the participants had an average attitude level in preventing oral and dental diseases. Also, 60% of the participants brushed their teeth only once a day, 22% brushed their teeth twice a day, and 8% did not brush at all. In addition, more than 66% of the participants did not floss their teeth to clean the area between their teeth. Previous studies have shown that only 22.9% of the participants brushed their teeth twice a day, and the majority (73.6%) did not floss their teeth [17, 29]. Given the important relationship between periodontal disease and diabetes, effective behaviors for preventing periodontal disease such as brushing, flossing, and regular dental visits not only can have positive impacts on the treatment of periodontal diseases, but they should be performed completely in order to maintain the health of diabetic patients [17].

The results of data analysis showed that 39% of the female patients visited a dentist for check-up, which is consistent with the results of previous studies [23-24, 30]. It was also revealed that 85% of the patients had not received any advice from a specific source to follow routine dental checkups. Physicians (7.4%) and friends (2.9%) were the main sources of suggestions for visiting a dentist. A study by Al Habashneh *et al.* [31] (2010) showed that for 50% of the patients, TV and Internet were the main cause of referral to dentists. In another study, only 5.2% of the patients were advised by their physicians to visit a dentist [17].

Studies have shown that patients who visited a dentist regularly, had a better understanding of the relationship between oral diseases and diabetes [17]. It is obvious that more knowledge of oral complications of diabe-

tes can help patients have a better attitude with respect to prevention and treatment of oral diseases. For this reason, public health education aims to improve patients' attitude via increasing their knowledge, and to help them take care of themselves. In recent studies conducted in Iran, it has been reported that the knowledge and attitude of diabetic patients about diabetes oral complications were poor, and none of the methods used in teaching patients about this problem was successful, hence, it seems that further studies should be done on this issue [31-32].

One of the potential reasons for the inadequate patients' knowledge may be the lack of awareness of the relationship between diabetes and periodontal disease among physicians and dentists. Hence, it is better to evaluate the awareness of health personnel regarding the relationship between diabetes and oral health in future studies. In addition, the information of medical and health personnel should be regularly updated [17].

Conclusion

According to the results, diabetic patients' knowledge and attitude of their oral and dental health is at moderate level. The analysis of the responses of the participants showed that female patients and those with longer diseases duration have a better knowledge and attitude of oral and dental health. Therefore, providing knowledge of periodontal and oral diseases to the public, especially diabetic patients, can be effective in improving the attitude in this area.

Conflict of Interest

The authors declare that they have no conflict of interests.

References

- [1] Pariookh M, Eghbal MJ, Ghodousi J, Kakoei S, Haghdoost AA, Kakoei S. The frequency of medically compromised patients in endodontic offices in iran. Iran Endod J. 2013; 8: 48-51.
- [2] Gambert SR. Are we up to the challenge? Clin Geriatr. 2002; 10: 14-26.
- [3] King H, Aubert RE, Herman WH. Global burden of diabetes, 1995-2025: prevalence, numerical estimates, and projections. Diabetes Care. 1998; 21: 1414-1431.
- [4] Smyth S, Heron A. Diabetes and obesity: the twin epid-

- emics. *Nat Med.* 2006; 12: 75-80.
- [5] Sadeghi MA, Kharazmi E, Javanbakht M, Alireza BM. Economics cost of diabetes in type II diabetic patients. *PAYESH.* 2012; 11: 443-449.
- [6] Khaledi S, Moridi G, Gharibi F. Survey of eight dimensions quality of life for patients with diabetes type II, referred to Sanandaj diabetes center in 2009. *J Fasa Univ Med Sci.* 2011; 1: 29-37.
- [7] Lankarani M, Zahedi F. Primary prevention of type 2 diabetes mellitus. *Iran J Diabetes Metab.* 2002; 1: 87-106.
- [8] Bharateesh J, Ahmed M, Kokila G. Diabetes and Oral Health: A Case-control Study. *Int J Prev Med.* 2012; 3: 806-809.
- [9] [9] Manfredi M, McCullough MJ, Vescovi P, Al-Kaarawi ZM, Porter SR. Update on diabetes mellitus and related oral diseases. *Oral Dis.* 2004; 10: 187-200.
- [10] Little JW, Falas DA, Miller C, Rhodus NL. *Dental Management of Medically compromised patient.* 9th ed. Missouri: Mosby; 2013. p. 280-288.
- [11] Darvish Moghaddam S, Moshtaghi Kashanian G, Hayatbakhsh M, Mehdipour A. Effect of pharmacologic doses of vitamin C on glycosylated hemoglobin in type 2 diabetic patients. *Iranian Journal of Endocrinology and Metabolism.* 2002; 4: 15-21.
- [12] Finney LS, Finney Mo, GanzalesCompoy. What the mouth has to say about diabetes. Careful examinations can overt serious complication. *Postgrad Med.* 1997; 102: 117-126.
- [13] Yuen HK, Onicescu G, Hill EG, Jenkins C. A survey of oral health education provided by certified diabetes educators. *Diabetes Res Clin Pract.* 2010; 88: 48-55.
- [14] American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care.* 2014; 37 Suppl 1: S81-S90.
- [15] Kakoei S, Hosseini B, Haghdoost AA, Sanjari M, Hashemipour MA, Gholamhosseinian A. The detection of salivary glucose, caries and periodontal status in diabetes mellitus patients. *J Oral Health Oral Epidemiol.* 2014; 3: 79-84.
- [16] Kakoei S, Hosseini B, Haghdoost AA, Sanjari M, Gholamhosseinian A, Afshar VF. Evaluation of Salivary Secretory Immunoglobulin A Levels in Diabetic Patients and Association with Oral and Dental Manifestations. *Sultan Qaboos Univ Med J.* 2015; 15: e507-e511.
- [17] Bahammam MA. Periodontal health and diabetes awareness among Saudi diabetes patients. *Patient Prefer Adherence.* 2015; 9: 225-233.
- [18] Heidary SH, Shirazi F, Sanjari M, Salimi S, Baljani E. Evaluation of effective factors on the patients with type 2 diabetes to institute endocrine & metabolism of Iran University of Medical Science. *Iranian Journal of Diabetes and Lipid Disorder.* 2010; 9: 365-375.
- [19] Yuen HK, Wolf BJ, Bandyopadhyay D, Magruder KM, Salinas CF, London SD. Oral health knowledge and behavior among adults with diabetes. *Diabetes Res Clin Pract.* 2009; 86: 239-246.
- [20] Delvarianzadeh M, Bagheri H, Sadeghian F. Effectiveness of diabetes dietary counseling on quality of life in type 2 diabetic patients. *IJDLD.* 2006; 5: 369-376.
- [21] Black M, Hawks J, Keene M. *Medical-surgical nursing clinical management for positive outcomes.* 6th ed. W.B Saunders Company. University of Nebraska at Omaha, United States; 2001. p. 12-13.
- [22] Wändell PE, Tovi J. The quality of life of elderly diabetic patients. *J Diabetes Complications.* 2000; 14: 25-30.
- [23] Eldarrat AH. Diabetic patients: their knowledge and perception of oral health. *Libyan J Med.* 2011; 6. doi: 10.3402/ljm.v6i0.5691.
- [24] Allen EM, Ziada HM, O'Halloran D, Clerehugh V, Allen PF. Attitudes, awareness and oral health-related quality of life in patients with diabetes. *J Oral Rehabil.* 2008; 35: 218-223.
- [25] Leite RS, Marlow NM, Fernandes JK, Hermayer K. Oral health and type 2 diabetes. *Am J Med Sci.* 2013; 345: 271-273.
- [26] Taiwo JO. Oral health education needs of diabetic patients in Ibadan. *Afr J Med Med Sci.* 2000; 29: 269-274.
- [27] Awartin F. Oral health knowledge and practices in Saudi diabetic female patients. *PODJ.* 2009; 29: 149-152.
- [28] Kakoei S, Navabi N, Aghaabbasi S, Hashemipour MA. Oral health related quality of life in patients with diabetes mellitus type 2 in the year 2012. *J Oral Health Oral Epidemiol.* 2016; 5: 186-191.
- [29] Stewart JE, Wager KA, Friedlander AH, Zadeh HH. The effect of periodontal treatment on glycemic control in patients with type 2 diabetes mellitus. *J Clin Periodontol.* 2001; 28: 306-310.
- [30] Bakhshandeh S, Murtooma H, Vehkalahti MM, Mofid R, Suomalainen K. Oral self-care and use of dental services among adults with diabetes mellitus. *Oral Health*

Prev Dent. 2008; 6: 279-286.

- [31] Al Habashneh R, Khader Y, Hammad MM, Almuradi M. Knowledge and awareness about diabetes and periodontal health among Jordanians. *J Diabetes Complications*. 2010; 24: 409-414.
- [32] Atarbashi Moghadam F, Haerian A, Sadat Salami M, Akhavan Karbasi MH, Fakhri-Tabatabayi R, Vaziri F.

Evaluation of knowledge, Attitude and Practice between Periodontal Disease and Diabetes. *JCHR*. 2013; 2: 124-130.

- [33] Taheri JB, Khalighi HR, Azimi S, Mortazavi H, Noormohammadi H, Tarahomi MR. Oral health Knowledge of Diabetic Patients before and after the Education Package. *AJDR*. 2012; 4: 47-52.