

Case Report

Mandibular Arteriovenous Malformation with Unusual Radiographic Appearance

Nafiseh Shamloo¹, DDS, MSc; Fatemeh Mashhadiabbas¹, DDS, MSc; Roohollah Safarpour², DDS;

¹ Dept. of Oral and Maxillofacial Pathology, Dental School, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

² Postgraduate Student, Dept. of Oral and Maxillofacial Pathology, Dental School, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

KEY WORDS

Arterio-venous malformation;
Mandible;
Treatment;

Received:

Revised:

Accepted:

ABSTRACT

Arteriovenous malformation (AVM) of head and neck is a rare vascular anomaly but when present, it is persistent and progressive in nature. It can also represent a lethal benign disease due to massive hemorrhage. There are several indications for treatment, including age, location, extension and type of vascular malformation. Endovascular therapy can effectively cure most lesions with limited tissue involvement. Surgery can be used in selected cases in combination with embolization. Here, we present a rare case of arteriovenous malformation of mandible with floating tooth appearance in a 11-year-old boy patient. Given the spectrum of imaging presentations seen and the crossover with other lesions, microscopic histopathological examination is the gold standard for diagnosis.

Corresponding Author: Safarpour R, Dept. of Oral and Maxillofacial Pathology, Dental School, Shahid Beheshti University of Medical Sciences, Tehran, Iran. Tel: +98-212240 3075(261) Email: fanoos1979@yahoo.com

Cite this article as:

Introduction

All vascular lesions until 1980's were referred to as hemangiomas. Classification of vascular lesions based on endothelial characteristics into hemangiomas and vascular malformation was done by Mullikin *et al.* [1] in 1982. In 1996 International Association of the Study of Vascular Abnormalities modified the classification as vascular tumors or vascular malformations (VMs) [2]. Vascular malformations based on blood flow are classified into lesions with low flow such as lymphatic, capillary, venous malformations also lesions with high-flow lesions such as arteriovenous malformations (AVMs) as well as arteriovenous fistulas [1]. Differentiation criteria between AVM and hemangioma are based on the clinical and histological characteristics. In general, there is no sign of hemangiomas at birth, they are characterized by a rapid growth phase with endothelial cell proliferation. Shortly after birth, hemangiomas grow fast usually faster than the child's growth, which then show a gradual involution. On the other hand, symptoms of AVMs are visible and they grow relative to the growth of the child and remain throughout the life. They characterized by the normal turnover rate of endothelial cell and di-

rect, permanent communication between arteries and veins [1,3]. As benign lesions, AVMs can arise in the any area of the body. In some rare cases, AVMs appear in hard tissue like bone/jaws and cause a highly unusual spectrum of symptoms [4-5]. Jaw intraosseous arteriovenous malformations have a high risk in life threatening hemorrhage either spontaneously or following oral surgeries [6-7]. In this paper, we report the case of mandibular intraosseous arteriovenous malformations in an 11-year-old boy with floating tooth appearance in radiographic feature.

Case Presentation

An 11-year boy with complaint of a swelling without pain of mandibular right posterior area and tooth mobility in that region for one month was referred to the Department of Oral and Maxillofacial Surgery, Shahid Beheshti University of Medical Sciences, Tehran, Iran. The patient did not report a history of trauma or medical problems. Clinical examination was normal. In the intraoral examination, a fluctuant swelling extending from first premolar to first molar with erythematous mucosa was seen. Cone beam computed tomography (CBCT) sh-



Figure 1: floating tooth appearance

owed an ill-defined unilocular radiolucency that extended from first molar to the distal of first premolar with floating tooth appearance (Figure 1). Bone destruction was seen in lingual and buccal crest which caused perforation in some areas. It seemed that mandibular canal was intact (Figure 2). On aspiration blood was seen. Due to clinical, radiographic and aspiration findings, central giant cell granuloma, leukemia and osteosarcoma were suggested as a differential diagnosis. Under local anesthesia, excisional biopsy of the lesion was done, it completely excised with two teeth and was submitted for the histologic examination. The gross was an irregular, creamy-brown soft tissue with elastic consistency, measuring 2.5 X 2 X 1 centimeters.

Histopathology sections showed a vascular lesion composed of a varying mixture of arteries, veins and associated small vessels within a fibrous background. Mixed inflammatory cells infiltration, giant cells, hemorrhage and curetted bone are seen (Figure 3a and b). The lesion is covered by keratinized epithelium which in some areas ulcerated and replaced by fibrinopurulent membrane (Figure 4). According to histopathologic and

radiographic findings, the diagnosis of arteriovenous malformation was done. There was no recurrence during a 6-month follow-up (Figure 5).

Discussion

The VMs are malformed communications which lined by inactive endothelium. While such conditions are generally congenital, they demonstrate an extra growth if disturbed by infections, trauma or endocrine fluctuations [8]. Deepa *et al.* [9] indicated that the average age of presentation of intraosseous arteriovenous malformations in jaws (j-AVMs) is 19 years. This case, there had no symptoms until 11 years when a tooth mobility was noticed. J-AVMs are very rare conditions but featured a higher tendency of life-threatening due to hemorrhage either spontaneously or due to surgery, therefore differential diagnosis must be taken into account in deciding about unexplained symptoms and pathology [4,10]. The prevalence of the complication in female is twice that of male, while in our case is male, with the highest incidence between the ages of 10 and 20 with extremes at 3 months and 74 years of age. The head




Figure 2: CBCT image; mandibular canal is intact




Figure 3a: Photomicrograph showing a mixture of thick-walled arteries and veins, **b:** along with capillary vessels. ($\times 200$ Magnification, H&E)




Figure 4: Photomicrograph showing an *ulcerated surface epithelium* ($\times 100$ Magnification, H&E)



Figure 5: Dental panoramic radiograph of the patient at follow up revealed obviously bone repair of alveolus

and neck region constitutes 50% of vascular lesions so that only a small number of cases happen in the jaws. In addition, the prevalence of the complication is twice in the mandible than maxilla, as we saw in this case [11]. AVMs have slow growth nature, for this reason, they are asymptomatic for long time and can be seen at any age [12]. They usually appear without symptoms bruit, swelling of soft tissues, dental loosening, altered color of skin and mucosa and paresthesia of the lower lip or chin [13,14]. Although this case has no history of dental visits, but the appearance of jaw swelling and tooth mobility has been observed suddenly during one month. Due to existence of teeth, complex morphology of jaw structures and vital structures in maxillofacial area, diagnosis and management of jaw AVMs are more difficult than other intrabony AVMs [6]. When AVM appears close to teeth, they create mobility and displacement, consistent with the slow-growing nature, like in the present case. Cases of root resorption have also been

Reported [15]. In our case, bone destruction is seen in buccal and lingual crest, noticeably erosion of the alveolar process with apparently floating teeth, but root resorption is not seen. Clinically, J-AVMs are seen following extractions, while brushing or biopsy procedures [8]. The radiographic appearances of these lesions are variable and can be similar to other lesions; therefore, having a definitive diagnosis based just on imaging is not easy [16]. Differential diagnosis of J-AVMs includes hemangioma, Langerhans's cell histiocytosis, simple bone cyst, Ewing's sarcoma, aneurysmal bone cyst, ameloblastoma, osteosarcoma, odontogenic myxoma, ameloblastic fibroma, central giant cell granuloma, odontogenic keratocyst, dentigerous, fibrous dysplasia and metastatic malignant tumors [8,17-18]. Despite of unilocular radiolucent lesion in our case, most commonly, these lesions show a multilocular radiolucent defect. The lesions can be large (soap bubble appearance) or small (honeycomb appearance). In addition, the lesions

may appear as an ill-defined radiolucent area or a well-defined, cystic-radiolucency. In the case of larger malformation, cortical expansion or a sunburst radiographic appearance are expected. It is possible to demonstrate the vascular nature of the lesions using angiography [19].

In few cases, there are notable osseous erosion of the alveolus with visible floating teeth [20], such as present case. These features can be seen in other lesions like ameloblastomas, osteosarcomas, myxomas and or fibrous dysplasia [21]. Biopsy and histopathological examination for precise diagnosis of AVMs is needed. Features of AVM histology are relatively similar, so that fibrous tissue and proliferative epithelial cells are usually found around the lesion and the remaining bony trabeculae are interspersed with endothelial cells and vascular structures [4,22], as the present case which the remaining bony trabeculae are interspersed with vascular structures (Figure 3b).

AVMs demonstrate a therapeutic challenge because of their hemodynamic characteristics and their modality of growth. Elimination of lesion, hemorrhage control during surgery and preventing of recurrence are main goals for treatment of AVMs. Radiotherapy may be only treatment option for inaccessible AVMs; still, any deformity of the bone as effects on the surrounding structures can be corrected and controlled using surgery [23]. Treatment also depends on size of lesion and degree of involvement of vital structures. Direct surgery, that can be accompanied with embolization of major afferent (feeding) vessels is the standard treatment. For lesions that require resection, radiographic embolization often is performed one to two days earlier than surgery process to minimize blood loss [19]. The present case, depending on the size of lesion, was also treated with direct surgery without embolization.

Through radical excision of the large lesion we can perform curative purpose and in general, reconstruction is carried out immediately using a tissue graft [24].

The patient has signed informed consent for the surgical procedure and necessary information for reporting this case.

Conclusion

Here, we presented a rare case of mandibular intraosseous AVM with floating tooth appearance in an 11-year-

old male patient. Taking into account the spectrum of imaging presentations observed and the crossover with other lesions, microscopic histopathological investigation is the “gold standard” for precise diagnosis.

Conflict of Interest

Not declare

References

- [1] Mulliken JB, Glowacki J. Hemangiomas and vascular malformations in infants and children: a classification based on endothelial characteristics. *Plast Reconstr Surg.* 1982; 69: 412-422.
- [2] Enjolras O, Mulliken JB. Vascular tumors and vascular malformations (new issues). *Adv Dermatol.* 1997; 13: 375-423.
- [3] Pawane P; Anshu, Gangane N. Hemangiomas versus arterio-venous malformations: role of elastic stains and mast cell density. *Indian J Pathol Microbiol.* 2014; 57: 191-195.
- [4] Blackhall KK, Ling E, Kunjur J. A rare case of an intraosseousarteriovenous malformation of the temporo-mandibular joint and mandible–Case report and literature review. *Int J Surg Case Rep.* 2020; 76: 394-398.
- [5] Rosen RJ, Nassiri N, Drury JE. Interventional management of high-flow vascular malformations. *Tech VascInterv Radiol.* 2013; 16: 22-38.
- [6] Li X, Su L, Wang D, Gui Z, Jiang M, Yang X, et al. Clinical and imaging features of intraosseous arteriovenous malformations in jaws: a 15-year experience of single centre. *Sci Rep.* 2020; 10:12046.
- [7] Starke RM, McCarthy D, Komotar RJ, Connolly ES. Somatic KRAS mutation found in sporadic arteriovenous malformations. *Neurosurgery.* 2018; 83: E14-E15.
- [8] Rudagi BM, Patil S, Hammannavar R, Jaiswal T. Mandibular Arteriovenous Malformation in an 8-year-old Child: A Case Report. *J Contemp Dent Pract.* 2016; 17: 85-89.
- [9] Deepa V, David CM, Lidiya A. Management of arteriovenous vascular malformation masquerading as a mucocele using sclerotherapy- review of literature and a case report. *Int J Pharm Sci Invent.* 2013; 2: 1-6.
- [10] Stapf C, Mast H, Sciacca RR, Berenstein A, Nelson PK, Gobin YP, et al. The New York Islands AVM Study: design, study progress, and initial results. *Stroke.* 2003; 34: e29-e33.

- [11] Patussi C, Sassi LM, Bixofis RB, Zanferrari FL, Targa R, Schussel JL. Vascular Malformation in the Mandible and Maxilla: An Interesting Case Series. *Int J Oral Maxillofac Pathol.* 2014; 5: 37-41.
- [12] Rosenberg TL, Suen JY, Richter GT. Arteriovenous malformations of the head and neck. *Otolaryngol Clin North Am.* 2018; 51: 185-195.
- [13] Spreafico R, Sordo L, Bellotto R, Schipano M, Rescaldani A, Parmigiani F. Arterio-venous malformation of the mandible. Case report and review of literature. *Acta Otorhinolaryngol Ital.* 2016; 36: 333-336.
- [14] Sakkas N, Schramm A, Metzger MC, Berlis A, Schmelzeisen R, Otten JE, et al. Arteriovenous malformation of the mandible: a life-threatening situation. *Ann Hematol.* 2007; 86: 409-413.
- [15] Bunel K, Sindet-Pedersen S. Central hemangioma of the mandible. *Oral Surg Oral Med Oral Pathol.* 1993; 75: 565-570.
- [16] Dhiman NK, Jaiswara C, Kumar N, Patne SC, Pandey A, Verma V. Central cavernous hemangioma of mandible: Case report and review of literature. *Natl J Maxillofac Surg.* 2015; 6: 209-213.
- [17] Dwivedi AN, Pandey A, Kumar I, Agarwal A. Mandibular arteriovenous malformation: A rare life-threatening condition depicted on multidetector CT angiography. *J Oral Maxillofac Pathol.* 2014; 18: 111-113.
- [18] Blackhall KK, Ling E, Kunjur J. A rare case of an intraosseous arteriovenous malformation of the temporomandibular joint and mandible - Case report and literature review. *Int J Surg Case Rep.* 2020; 76:394-398.
- [19] Neville BW, Damm DD, Allen CM, Bouquot JE. *Oral and Maxillofacial Pathology.* 4th ed. St. Louis: Elsevier; 2016. p. 506,508.
- [20] Dwivedi AN, Pandey A, Kumar I, Agarwal A. Mandibular arteriovenous malformation: A rare life-threatening condition depicted on multidetector CT angiography. *J Oral Maxillofac Pathol.* 2014; 18:111-113.
- [21] Zlotogorski A, Buchner A, Kaffe I, Schwartz-Arad D. Radiological features of central haemangioma of the jaws. *Dentomaxillofac Radiol.* 2005; 34: 292-296.
- [22] Thoma KH, Goldman HM. Bone tumors of the jaws. *Thoma's Oral Pathology.* 1971; 6: 564-566.
- [23] Bhandari PS, Sadhotra LP, Bhargava P, Bath AS, Mukherjee MK, Maurya S. Management strategy for facial arteriovenous malformations. *Indian J Plast Surg.* 2008; 41: 183-189.
- [24] Kaneko R, Tohnai I, Ueda M, Negoro M, Yoshida J, Yamada Y. Curative treatment of central hemangioma in the mandible by direct puncture and embolisation with n-butyl-cyanoacrylate (NBCA). *Oral Oncol.* 2001; 37: 605-608.