

Original Article**Pain Perception Due to Dental Injection by Smartject: Split Mouth Design Study**Faezeh Ghaderi ¹, Mahboobe Ahmadbeigi ²¹ Oral and Dental Disease Research Center, Dept. of Pediatric Dentistry, School of Dentistry, Shiraz University of Medical Sciences, Shiraz, Iran.² Post Graduate Dental Student, Member of Research Committee, Shiraz University of Medical Sciences, Shiraz, Iran.**KEY WORDS**Anesthesia, Local;
Pain Perception;
Injections;
Visual Analogue Scale;**ABSTRACT****Statement of the Problem:** Dental injection is one of the most fearful procedures in dental setting, especially for children. Many researchers have attempted to find a painless method. As computer controlled local anesthesia delivery system devices (CCLADs) allow the speed rate and pressure of injection solution to be controlled, they may cause less pain during injection in comparison to the conventional method.**Purpose:** The aim of this study was to compare pain perception in dental injection by Smartject with conventional technique.**Materials and Method:** The present study was a randomized single-blind crossover clinical trial. The participants consisted of 50 healthy volunteer dental students.They received a topical anesthetic agent plus injection in maxillary premolar buccal mucosa via conventional technique on one side (control) and a topical anesthetic agent plus injection in maxillary premolar buccal mucosa by Smartject on the other side (experimental). The first injection method was chosen based on block randomization table. A blind person recorded the subjects' pain perception of injection based on the visual analogue scale (VAS) in the two groups. Repeated measure test, independent Student t-test and Student paired t- test were used. Statistical significance was defined at $p < 0.05$.**Results:** There was statistically significant difference in VAS score between Smartject and the conventional technique. The mean of VAS scores for Smartject and the conventional technique were 14.5 ± 7.4 and 24 ± 12.1 , respectively.**Conclusion:** It is suggested, needle penetration is not the main reason of pain during injection. Inconsistent fluid pressure created by injected anesthetic solution on nerve fibers is more impressive in pain development. Hence, Smartject as a CCLAD can be considered as an appropriate device for dental injection.Received May 2017;
Received in Revised form June 2017;
Accepted July 2017;**Corresponding Author:** Ahmadbeigi M., Ghasrodasht St, Dental School of Shiraz University of Medical Sciences, Shiraz, Iran. Email: mahboobe.ahmadbeigi@gmail.com Fax: +98-7136270325
Tel: +98-7136263193-4Cite this article as: Ghaderi F., Ahmadbeigi M. Pain Perception Due to Dental Injection by Smartject: Split Mouth Design Study. *J Dent Shiraz Univ Med Sci.* 2018 March; 19(1): 57-62.**Introduction**

According to many studies, the most fearful experience cited by subjects who were asked to rank the situations in dental treatment is fear of dental injection. [1-2] This fear may prevent patients from seeking dental treatment. [3]

Numerous studies have been conducted to find a painless injection, [4-6] including application of topical

anesthesia, [7] prolonged injection time, [8] pre-cooling injection site, [9] pressure to injection site, [10] applying laser as pretreatment method, [11-13] warming the local anesthetic agents, [14] buffering the local anesthetics, [14-15] tactile stimulation, [16-17] and distraction technique. [18] However, more situations and techniques need to be evaluated to improve stress management in dental setting.

Traditional injection systems that uses metallic syringe do not allow the control of flow rate in a constant way and consequently, fluid pressure varies depending on manual force used by the practitioner during the injection procedure. [19] An injection into some areas in oral cavity, such as the palate that requires harder force, causes difficulty in syringe control and does not provide a comfortable injection. [6]

Recently, several researches have focused on the effectiveness of computer controlled local Anesthesia delivery system devices (CCLADs). These devices are designed to control the pain of the injection by delivering anesthetic solution at a constant pressure, volume and injection rate. [19] Several companies have improved the computer based, speed-controlled local anesthetic devices, including Comfort Control Syringe (CCS®) and the Comfort-in®, Deninjection®, No Pain III®, Meg-inject®, and Smartject®. These devices have different characteristics, such as design, weight, injection speed, shape, and possibility of aspiration, so operator can choose the device based on the convenience, depending on different situations. [20]

Smartject has been developed to provide operator convenience. Its lightweight design provides safe injection; in heavy design devices, it is difficult to control movement during needle injection, especially when a long period is needed for injection. [20] Moreover, having the possibility to use standard lidocaine carpule as well as conventional dental needle are the other advantages of Smartject. [20] Another good point of Smartject is having the possibility of aspiration; it is a facility where block injection is needed. [6]

Many studies have shown anxiety as well as pain perception in pediatric and general population could be reduced with CCLAD system. [21-27] However, some studies found no statistically significant difference in pain perception between conventional injection and a computerized device. [19, 22, 28]

It is worth mentioning, that so far no other studies have compared Smartject with the conventional technique. In addition, there is a controversial result about CCLAD efficacy in comparison with the conventional technique. [21, 28-29] The current study was designed to evaluate Smartject efficacy in comparison with conventional syringe injections on pain related with dental injection.

Materials and Method

The current study was a randomized single-blind cross-over clinical trial and it was conducted in Dental School of Shiraz University of Medical Sciences, Shiraz, Iran. Fifty healthy volunteer dental school students from Shiraz University of Medical Sciences participated in this study. Their age ranged from 23 to 28 years. The sample size was determined based on a similar study. [30]

The exclusion criteria were history of systemic disease, presence of abscess, redness, fistula in injection site, taking any kinds of medication that would change the pain perception, and history of allergy to lidocaine. The inclusion criteria were considered as all participants should have previously experienced the conventional syringe and none of them had ever experienced Smartject injection before.

After explaining the study procedure, a written consent form was obtained from all participants before the commencement of the trial. The Committee of Research and Ethics of university approved all the aspects and steps of this research protocol (Grant#94-01-03-9946) and it was submitted in IRCT (IRCT2015101024-445N2).

The participants received a topical anesthetic agent followed by injection by conventional technique on one side (control) and a topical anesthetic agent followed by injection by Smartject on the other side (experimental). The first injection method was chosen based on block randomization table.

The patients were explained how to report their pain perception during injection using visual analogue scale (VAS). VAS was scored on a 100-mm horizontal line with the left endpoint marked “no pain” and the right endpoint marked “pain as bad as it can be” for each patient.

Injection procedure

Buccal infiltration injection was conducted in maxillary premolar buccal mucosa aiming for the apical area of the tooth to evaluate the pain perception. On the experimental side, the injection site was dried with a cotton roll for thirty seconds and then benzocaine gel (Benzocaine USP, USA) was applied for one minute. The procedure was immediately followed by injection with mode 1 Smartject (KMG co, Saha-gu, Busan, Korea) (Figure 1) using 30 gauge short needle (DENJECT, Korea). In the same session, on the control side, injec-

tion site was dried with a cotton roll for thirty seconds and then Benzocaine gel (Benzocaine USP, USA) was applied for one minute. The procedure was immediately followed by conventional injection with 30 gauge short needles (DENJECT, Korea). In both the experimental and control sides, 1 ml of 2% lidocaine with 1:80,000 epinephrine (Lidocaine 2% E-80, Colombia) was injected. A trained examiner, blinded to the study, recorded pain perception during injection in both sides based on VAS.



Figure 1: Smartject

Data were analyzed by SPSS software version 21. Repeated measure test was used to find out the effect of treatment sequence on results. Student paired t- test was applied to compare VAS score between Smartject and the conventional technique, Independent Student t-test was used to compare VAS score between male and female in the conventional technique as well as Smartject. Statistical significance was defined at $p < 0.05$.

Results

In this study, the total number of participant was 50 and male to female ratio was 22 males (44%) and 28 females (56%), their age ranged from 23 to 28 (24.3 ± 1.7) years. Repeated measure test showed that the sequence of treatment had no effect on the result. Hence, in this study we merely evaluated the pain perception of the two mentioned methods. Student paired t- test showed statistically significant difference between the mean of VAS scores for Smartject and the conventional technique (14.5 ± 7.4 and 24 ± 12.1 respectively). ($p = 0.006$)

Student paired t- test showed statistically significant difference in the VAS score between Smartject and the conventional technique in males as well as females (Table 1).

Table 1: Comparison of Visual Analogue Scale in Smartject and conventional technique based on the gender

Gender	Number	Mean±SD VAS (Smartject)	Mean±SD VAS (conventional technique)	p Value
Male	22	12.5±8.07	25.3±11.3	0.000
Female	28	15.1±10.2	22.2±10.1	0.011

Independent Student t-test showed no statistically significant difference in VAS score between males and females in the conventional technique as well as Smartject (Table 2).

Table 2: Comparison of Visual Analogue Scale in male and female in both injection methods

Injection method	Mean±SD VAS in Male	Mean±SD VAS in Female	p Value
Smartject	12.5±8.07	15.1±10.2	0.420
Conventional technique	25.3±11.3	22.2±10.1	0.450

Discussion

Results of the current study showed the buccal infiltration injection by Smartject device caused statistically less pain in comparison with the conventional syringe in male and female, which was clinically important (more than five scores in VAS). [30] Different studies reported a range of at least 13-30 mm decrement in VAS score as clinically significant. [31-32]

Several studies showed anterior middle superior alveolar injection with computer controlled devices to be less painful than that using conventional syringe technique. [21, 33-34]

Krochak and Friedman stated that patients were able to manage their anxiety about injections successfully after applying the Wand system (a type of CCLAD). Hence, this can be a good method to decrease disruptive behaviors in children, especially in younger children. [35] Moreover, children who had experienced conventional technique were five times more likely needing be controlled for dental management than children who received injection by a computer controlled devices. [19]

Since CCLADs provide precise control of anesthetic solution flow rate, pressure and duration of injection time, the factors that are so difficult to achieve by manual injection, these results could be expected. However, some studies reported that there is no difference in pain perception during injection with conventional syringe in comparison to CCLAD devices. [22, 28-29]

As pain is a multi-factorial issue, it can be influenced by different factors such as age, gender, and situation. [36] The wide age range in the two above-mentioned studies [28-29] should be considered as their limitation for the same reason. The age range of participants in Versloot *et al.* [28] study and in Asarch *et al.* [29] study was 4-11 years and 5-13 years, respectively.

Allen *et al.* [19] reported after the first 15-second interval of the injection by computerized injection device, disruptive behavior developed as the time passed. In contrast to Allen *et al.* [19] study, Gibson *et al.* [37] study reported disruptive behavior diminished over time. However, in current study this parameter was not assessed, as the subjects were adult. In common belief, needle penetration into the tissue is not the main reason contributing to discomfort. Volume and pressure induced by anesthetic solution during injection results in more pain and stress. Anxiety in patients following local anesthetic injection makes the dentist nervous. [5, 21]

In current study, gender was not found to be a significant factor in pain perception in both injection methods. This result was in agreement with some other studies. [17, 38] However, in some studies gender was reported as a factor in pain perception as well as in acceptance of new method. [11, 16, 39] This discrepancy could be due to different inclusion criteria such as age and awareness level about applied procedure.

Split mouth design of this study provided results that are more reliable. The subjects for this study were adult because perception of pain in preschool children is complex and involves behavioral, physiological, psychological, and developmental factors. [19] The single blind design of the study can be considered as a limitation. In addition, limited number of Smartject cartridges (where carpules and needle are mounted) resulted in taking more time. The fact that the participants were drawn from dental students might give rise to possible bias. Evaluating different age groups, as well as different type of injection and other CCLAD devices are recommended for future studies.

Conclusion

It seems that needle insertion might not be the most important factor on pain perception during an injection. It seems that inconsistent fluid pressure and speed created by anesthetic solution on the nerve fibers is more significant in pain perception during conventional injection method. Therefore, Smartject (as a CCLAD) can be considered as an appropriate device for dental injection owing to its several options and aptitudes.

Acknowledgments

The authors would like to thank the Vice-Chancellor of

Research at Shiraz University of Medical Sciences for supporting this research (Grant # 94-01-03-9946). This article is based on a thesis by Dr. Mahboobe Ahmad-beigi. The authors also thank Dr. Salehi of the Dental Research Development Center, School of Dentistry for the statistical analysis and would like to acknowledge Research Deputy of Shiraz Dental School. The authors wish to thank the Research Consultation Center (RCC) at Shiraz University of Medical Sciences for their invaluable assistance in editing this manuscript.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- [1] Milgrom P, Coldwell SE, Getz T, Weinstein P, Ramsay DS. Four dimensions of fear of dental injections. *J Am Dent Assoc.* 1997; 128: 756-766.
- [2] Ghaderi F, Banakar S, Rostami S. Effect of pre-cooling injection site on pain perception in pediatric dentistry: "A randomized clinical trial". *Dent Res J (Isfahan).* 2013; 10: 790-794.
- [3] Larsson BA, Gradin M, Lind V, Selander B, Swedish Paediatric Pain Society. Swedish guidelines for prevention and treatment of pain in the newborn infant. *Lakar-tidningen.* 2002; 99: 1946-1949.
- [4] Carbajal R, Lenclen R, Gajdos V, Jugie M, Paupe A. Crossover trial of analgesic efficacy of glucose and pacifier in very preterm neonates during subcutaneous injections. *Pediatrics.* 2002; 110(2 Pt 1): 389-393.
- [5] Simon JF, Peltier B, Chambers D, Dower J. Dentists troubled by the administration of anesthetic injections: long-term stresses and effects. *Quintessence Int.* 1994; 25: 641-646.
- [6] Salgotra V, Agrawal Rm Mandal S, Kohli S. New Gadgets in Local Anesthesia: A review. *IOSR-JDMS.* 2014; 13: 62-66.
- [7] O'Brien L, Taddio A, Lyszkiewicz DA, Koren G. A critical review of the topical local anesthetic amethocaine (Ametop) for pediatric pain. *Paediatr Drugs.* 2005; 7: 41-54.
- [8] Courtney DJ, Agrawal S, Revington PJ. Local anaesthesia: to warm or alter the pH? A survey of current practice. *J R Coll Surg Edinb.* 1999; 44: 167-171.
- [9] Aminabadi NA, Farahani RM. The effect of pre-cooling

- the injection site on pediatric pain perception during the administration of local anesthesia. *J Contemp Dent Pract.* 2009; 10: 43-50.
- [10] Johnson J, Primosch RE. Influence of site preparation methods on the pain reported during palatal infiltration using the Wand Local Anesthetic System. *Am J Dent.* 2003; 16: 165-169.
- [11] Ghaderi F, Ghaderi R, Davarmanesh M, Bayani M, Arabzade Moghadam S. Pain management during needle insertion with low level laser. *Eur J Paediatr Dent.* 2016; 17: 151-154.
- [12] Armida MMT. Laser therapy and its applications in dentistry. *Pract Odontol.* 1989; 10: 9-16.
- [13] Shapiro H, Harris L, Hetzel FW, Bar-Or D. Laser assisted delivery of topical anesthesia for intramuscular needle insertion in adults. *Lasers Surg Med.* 2002; 31: 252-256.
- [14] Colaric KB, Overton DT, Moore K. Pain reduction in lidocaine administration through buffering and warming. *Am J Emerg Med.* 1998; 16: 353-356.
- [15] Younis I, Bhutiani RP. Taking the 'ouch' out- effect of buffering commercial xylocaine on infiltration and procedure pain- a prospective, randomised, double-blind, controlled trial. *Ann R Coll Surg Engl.* 2004; 86: 213-217.
- [16] Hogan ME, Probst J, Wong K, Riddell RP, Katz J, Taddio A. A randomized-controlled trial of parentled tactile stimulation to reduce pain during infant immunization injections. *Clin J Pain.* 2014; 30: 259-265.
- [17] Sparks L. Taking the "ouch" out of injections for children. Using distraction to decrease pain. *MCN Am J Matern Child Nurs.* 2001; 26: 72-78.
- [18] Touyz LZ, Lamontagne P, Smith BE. Pain and anxiety reduction using a manual stimulation distraction device-when administering local analgesia oro-dental injections: a multi-center clinical investigation. *J Clin Dent.* 2004; 15: 88-92.
- [19] Allen KD, Kotil D, Larzelere RE, Hutfless S, Beiraghi S. Comparison of a computerized anesthesia device with a traditional syringe in preschool children. *Pediatr Dent.* 2002; 24: 315-320.
- [20] Kwak EJ, Pang NS, Cho JH, Jung BY, Kim KD, Park W. Computer-controlled local anesthetic delivery for painless anesthesia: a literature review. *J Dent Anesth Pain Med.* 2016; 16: 81-88.
- [21] Yenisey M. Comparison of the pain levels of computer-controlled and conventional anesthesia techniques in prosthodontic treatment. *J Appl Oral Sci.* 2009; 17: 414-420.
- [22] Ram D, Peretz B. The assessment of pain sensation during local anesthesia using a computerized local anesthesia (Wand) and a conventional syringe. *J Dent Child (Chic).* 2003; 70: 130-133.
- [23] Palm AM, Kirkegaard U, Poulsen S. The wand versus traditional injection for mandibular nerve block in children and adolescents: perceived pain and time of onset. *Pediatr Dent.* 2004; 26: 481-484.
- [24] Singh S, Garg A. Comparison of the pain levels of computer controlled and conventional anesthesia techniques in supraperiosteal injections: a randomized controlled clinical trial. *Acta Odontol Scand.* 2013; 71: 740-743.
- [25] Nicholson JW, Berry TG, Summitt JB, Yuan CH, Witten TM. Pain perception and utility: a comparison of the syringe and computerized local injection techniques. *Gen Dent.* 2001; 4: 167-173.
- [26] Fukayama H, Yoshikawa F, Kohase H, Umino M, Suzuki N. Efficacy of anterior and middle superior alveolar (AMSA) anesthesia using a new injection system: the Wand. *Quintessence Int.* 2003; 34: 537-541.
- [27] Perry DA, Loomer PM. Maximizing Pain Control. The AMSA Injection can provide anesthesia with few injections and less pain. *Dimens Dent Hyg.* 2003; 1: 28-33.
- [28] Versloot J, Veerkamp JS, Hoogstraten J. Pain behaviour and distress in children during two sequential dental visits: comparing a computerised anaesthesia delivery system and a traditional syringe. *Br Dent J.* 2008; 205: E2;discussion 30-31.
- [29] Asarch T, Allen K, Petersen B, Beiraghi S. Efficacy of a computerized local anesthesia device in pediatric dentistry. *Pediatr Dent.* 1999; 21: 421-424.
- [30] Lee JS. Pain measurement: understanding existing tools and their application in the emergency department. *Emerg Med (Fremantle).* 2001; 13: 279-287.
- [31] Gallagher EJ, Liebman M, Bijur PE. Prospective validation of clinically important changes in pain severity measured on a visual analog scale. *Ann Emerg Med.* 2001; 38: 633-638.
- [32] Lee JS, Hobden E, Stiell IG, Wells GA. Clinically important change in the visual analog scale after adequate pain control. *Acad Emerg Med.* 2003; 10: 1128-1130.
- [33] Loomer PM, Perry DA. Computer-controlled delivery versus syringe delivery of local anesthetic injections for therapeutic scaling and root planning. *J Am Dent Assoc.* 2004; 135: 358-365.
- [34] Ashwini S, Bhavya B, Nambiar M, Shireen S. Comparis-

- on of Pain Intensity of AMSA Injection with Conventional Syringe and WAND System for Maxillary Periodontal Surgery. *International Journal of Advanced Research*. 2014; 2: 349-353.
- [35] Krochak M, Friedman N. Using a precision metered injection system to minimize dental injection anxiety. *Compendium*. 1998; 19: 137-148.
- [36] Wandner LD, Scipio CD, Hirsh AT, Torres CA, Robinson ME. The perception of pain in others: how gender, race, and age influence pain expectations. *J Pain*. 2012; 13: 220-227.
- [37] Gibson RS, Allen K, Hutfless S, Beiraghi S. The Wand vs. traditional injection: a comparison of pain related behaviors. *Pediatr Dent*. 2000; 22: 458-462.
- [38] Jones A, Zachariae R. Investigation of the interactive effects of gender and psychological factors on pain response. *Br J Health Psychol*. 2004; 9(Pt 3): 405-418.
- [39] Craft RM, Mogil JS, Aloisi AM. Sex differences in pain and analgesia: the role of gonadal hormones. *Eur J Pain*. 2004; 8: 397-411.